NASA/TM-2000-209891, Vol. 34



Technical Report Series on the Boreal Ecosystem-Atmosphere Study (BOREAS)

Forrest G. Hall and David E. Knapp, Editors

Volume 34 BOREAS HYD-8 1994 Gravimetric Moss Moisture Data

X. Wang

National Aeronautics and Space Administration

Goddard Space Flight Center Greenbelt, Maryland 20771

The NASA STI Program Office ... in Profile

Since its founding, NASA has been dedicated to the advancement of aeronautics and space science. The NASA Scientific and Technical Information (STI) Program Office plays a key part in helping NASA maintain this important role.

The NASA STI Program Office is operated by Langley Research Center, the lead center for NASA's scientific and technical information. The NASA STI Program Office provides access to the NASA STI Database, the largest collection of aeronautical and space science STI in the world. The Program Office is also NASA's institutional mechanism for disseminating the results of its research and development activities. These results are published by NASA in the NASA STI Report Series, which includes the following report types:

- TECHNICAL PUBLICATION. Reports of completed research or a major significant phase of research that present the results of NASA programs and include extensive data or theoretical analysis. Includes compilations of significant scientific and technical data and information deemed to be of continuing reference value. NASA's counterpart of peer-reviewed formal professional papers but has less stringent limitations on manuscript length and extent of graphic presentations.
- TECHNICAL MEMORANDUM. Scientific and technical findings that are preliminary or of specialized interest, e.g., quick release reports, working papers, and bibliographies that contain minimal annotation. Does not contain extensive analysis.
- CONTRACTOR REPORT. Scientific and technical findings by NASA-sponsored contractors and grantees.

- CONFERENCE PUBLICATION. Collected papers from scientific and technical conferences, symposia, seminars, or other meetings sponsored or cosponsored by NASA.
- SPECIAL PUBLICATION. Scientific, technical, or historical information from NASA programs, projects, and mission, often concerned with subjects having substantial public interest.
- TECHNICAL TRANSLATION.
 English-language translations of foreign scientific and technical material pertinent to NASA's mission.

Specialized services that complement the STI Program Office's diverse offerings include creating custom thesauri, building customized databases, organizing and publishing research results . . . even providing videos.

For more information about the NASA STI Program Office, see the following:

- Access the NASA STI Program Home Page at http://www.sti.nasa.gov/STI-homepage.html
- E-mail your question via the Internet to help@sti.nasa.gov
- Fax your question to the NASA Access Help Desk at (301) 621-0134
- Telephone the NASA Access Help Desk at (301) 621-0390
- Write to:
 NASA Access Help Desk
 NASA Center for AeroSpace Information
 7121 Standard Drive
 Hanover, MD 21076-1320

NASA/TM-2000-209891, Vol. 34



Technical Report Series on the Boreal Ecosystem-Atmosphere Study (BOREAS)

Forrest G. Hall and David E. Knapp, Editors

Volume 34 BOREAS HYD-8 1994 Gravimetric Moss Moisture Data

Xuewen Wang, University of Toronto

National Aeronautics and Space Administration

Goddard Space Flight Center Greenbelt, Maryland 20771

			•	
		-		
		•		•
	•	Available from	ı:	•
NASA Center for AeroSpace I 7121 Standard Drive Hanover, MD 21076-1320 Price Code: A17	nformation		National Te	chnical Information Serv 5285 Port Royal Ro Springfield, VA 221 Price Code: A

BOREAS HYD-8 1994 Gravimetric Moss Moisture Data

Xuewen Wang

Summary

The BOREAS HYD-8 team made measurements of surface hydrological processes that were collected at the NSA-OBS Tower Flux site in 1994 and at Joey Lake, Manitoba, to support its research into point hydrological processes and the spatial variation of these processes. The data collected may be useful in characterizing canopy interception, drip, throughfall, moss interception, drainage, evaporation, and capacity during the growing season at daily temporal resolution. This particular data set contains the gravimetric moss moisture measurements from June to September 1994. A nested spatial sampling plan was implemented to support research into spatial variations of the measured hydrological processes and ultimately the impact of these variations on modeled carbon and water budgets. These data are stored in tabular ASCII files.

Table of Contents

- 1) Data Set Overview
- 2) Investigator(s)
- 3) Theory of Measurements
- 4) Equipment
- 5) Data Acquisition Methods
- 6) Observations
- 7) Data Description
- 8) Data Organization
- 9) Data Manipulations
- 10) Errors
- 11) Notes
- 12) Application of the Data Set
- 13) Future Modifications and Plans
- 14) Software
- 15) Data Access
- 16) Output Products and Availability
- 17) References
- 18) Glossary of Terms
- 19) List of Acronyms
- 20) Document Information

1. Data Set Overview

1.1 Data Set Identification

BOREAS HYD-08 1994 Gravimetric Moss Moisture Data

1.2 Data Set Introduction

This particular data set contains the gravimetric moss moisture measurements by the Hydrology (HYD)-08 team at the BOReal Ecosystem-Atmosphere Study (BOREAS) Northern Study Area (NSA)-Old Black Spruce (OBS) site and the Joey Lake, Manitoba, site from June to September 1994. A nested spatial sampling plan was implemented to support research into spatial variations of the measured hydrological processes and ultimately the impact of these variations on modeled carbon and water budgets. These data are stored in American Standard Code for Information Interchange (ASCII) text files.

1.3 Objective/Purpose

The objective of these moss data is to provide the BOREAS investigators with a data product that characterizes the role of moss on the water budget at the forest floor in the NSA-Modeling Sub-Area (MSA). These data are to be used for modeling purposes.

1.4 Summary of Parameters

This data set contains the weight of moss turfs for different layers of moss under conditions where the turfs were contained in plastic, to contain water within the moss turf, and without plastic, to let water flow through the moss.

1.5 Discussion

Hydrological processes such as canopy evaporation, moss storage, and moss evaporation may play a significant role in controlling water fluxes during the growing season in boreal wetlands. Canopy interception, moss storage, and moss evaporation were measured using mass balance methods (throughfall catch buckets and lysimeters) to give a quantitative estimate of these processes for sparse black spruce stands. More importantly, the spatial sampling scheme allowed quantification of the expected variation of these processes within the footprint of a colocated flux measurement tower. This will allow consideration of the subtower-footprint controls on vapor fluxes that the tower is measuring. In addition, the data set will be useful in parameterizing flux models for the targeted site, as well as determining the typical variation in fine-scale processes that the models may have to account for when scaling to watershed and regional extents.

1.6 Related Data Sets

BOREAS HYD-01 Volumetric Soil Moisture Data BOREAS HYD-01 Under Canopy Precipitation BOREAS HYD-06 Moss/Humus Moisture Data BOREAS HYD-08 Throughfall Data BOREAS HYD-08 1996 Gravimetric Moss Moisture Data BOREAS HYD-08 Gross Precipitation Data

2. Investigator(s)

2.1 Investigator(s) Name and Title

Dr. Lawrence Band University of North Carolina Chapel Hill, NC

Formerly at: University of Toronto Department of Geography Toronto, Ontario

2.2 Title of Investigation

Simulation of Boreal Ecosystem Carbon and Water Budgets: Scaling from Local to Regional Extents

2.3 Contact Information

Contact 1:

Xuewen Wang University of Toronto Department of Geography Toronto, Ontario M5T 1A1 Canada (416) 978-5070 wangx@esker.geog.utoronto.ca

Contact 2:

L.E. Band University of Toronto Department of Geography Toronto, Ontario M5T 1A1 Canada (416) 978-3375 lband@eos.geog.utoronto.ca

Contact 3:

David Knapp
Raytheon ITSS
NASA GSFC
Code 923
Greenbelt, MD 20771
(301) 286-1424
(301) 286-0239 (fax)
David.Knapp@gsfc.nasa.gov

3. Theory of Measurements

Turfs were located randomly; however, some turfs were relocated because of the presence of live roots. A set of 20-cm x 20-cm-square turfs was extracted by hand either to the bottom of the live moss layer (defined by the presence of a horizontal litter mat) for Pleurozium schreberi turfs, or by the end of thalli for sphagnum turfs. Turfs were placed on a mesh tray and then replaced in the pit from which they were extracted. Turfs were weighed daily and after rain events by placing them directly on a leveled electronic scale. If turfs were over field capacity (water dripped from them), the phenomenon was noted and weighing proceeded after the majority of drip water had ceased. Any large litter components that appeared on turfs during the measurement period were removed and placed on the moss surface beside the turf. Turf water equivalent depth was computed by later oven drying the turfs and determining dry turf + lysimeter tray weights to compute water weight, assuming a constant 1,000 kg/m³ density of H_2O to compute water equivalent depth.

4. Equipment

4.1 Sensor/Instrument Description

Turf Lysimeter Mesh Base

A 20-cm x 20-cm aluminum mesh tray with 10-gauge wire mesh and 1-cm mesh holes was used as the base of each lysimeter. Monofilament fishing line was used to form handles to extract the lysimeter

from the turf pit.

Electronic Weigh Scales

The two scales used were manufactured by MARS, and were both in the MS3000W series. The scale used to measure weights less than 1 kg was a 900-g range scale (accurate to \pm 0.1 g). The scale used to measure weights greater than 1 kg was a 2,000-g range scale (accurate to \pm 1.0 g). Both balances were tared before and after weighing and had been calibrated immediately before the field campaign and at the University of Toronto after the measurement campaign. The balances had an auto-off condition where taring was not possible or when battery power was low.

4.1.1 Collection Environment

These measurements were made in a spruce forest, near the NSA-OBS tower site and the Joey Lake site, where moss covered the ground.

4.1.2 Source/Platform

None.

4.1.3 Source/Platform Mission Objectives

None.

4.1.4 Key Variables

Moss turf weights.

4.1.5 Principles of Operation

See Section 3.

4.1.6 Sensor/Instrument Measurement Geometry

All throughfall and catch gauges were repositioned using a bubble level to ensure that they were upright. The turf lysimeters were extracted vertically from the pits even for pits on the sides of hummocks.

4.1.7 Manufacturer of Sensor/Instrument

Gauges and Lysimeters - Darryl Carlysle Moses and Kira Dunham (University of Toronto, Dept. of Geography)

Weigh Scales - (2) MARS MS3000W Series

4.2 Calibration

The weigh scales were calibrated to within the manufacturer's specifications immediately before the measurement campaign and at the University of Toronto after the campaign. The effect of the weigh scales being off level was also tested with no appreciable difference for tilt angles less than 20 degrees (which were defined by the first indent in the bubble level gauge used in the field).

4.2.1 Specifications

None given.

4.2.1.1 Tolerance

None given.

4.2.2 Frequency of Calibration

The weigh scales were calibrated to within the manufacturer's specifications immediately before the measurement campaign and at the University of Toronto after the campaign.

4.2.3 Other Calibration Information

None.

5. Data Acquisition Methods

The living moss carpet was cut in 20-cm by 20-cm squares with varying depths: L layer consists of the living moss carpet and a small portion of the brown senescing moss shoot bases; LF layer includes L layer plus a brown, well-defined layer of partially decomposed but still recognizable organic material; and LFH layer is LF plus the very dark layer of more humified organic material above the Ae horizon of the mineral soil. Two treatments were applied, one with a plastic sheet at the bottom to prevent drainage from the turf and the other without plastic. The cut turfs were placed back in their original holes and taken out for weighing on a regular basis.

6. Observations

6.1 Data Notes

None given.

6.2 Field Notes

None given.

7. Data Description

7.1 Spatial Characteristics

7.1.1 Spatial Coverage

At the Joey Lake site, these data were collected at four different plots. At the NSA-OBS site, the data were taken from two plots located near the flux tower. The plots were about 30 meters apart, about 10 m by 10 m square each. The samples within each plot were about 5 meters apart. Each turf sample was 20 cm by 20 cm in size.

The approximate locations of the various plots (based on the North American Datum of 1983

(NAD83)) are as follows:

		•	BOREAS	Grid	
Site	Longitude	Latitude	X	Y	
NSA (Joey Lake)	98.15026W	55.46676N	807.025	571.969	
NSA-OBS (Flux Twr.)	98.48139W	55.88007N	778.216	613.516	

7.1.2 Spatial Coverage Map

None.

7.1.3 Spatial Resolution

The samples were 20 cm by 20 cm, randomly distributed around the given locations.

7.1.4 Projection

These data were collected at point locations. They are not in any projection.

7.1.5 Grid Description

None.

7.2 Temporal Characteristics

7.2.1 Temporal Coverage

NSA-Joey Lake: 25-Jun-1994 to 07-Sep-1994 NSA-OBS: 15-Aug-1994 to 16-Sep-1994

7.2.2 Temporal Coverage Map

None.

7.2.3 Temporal Resolution

These data were collected once a day (twice a day when it rained) for most days during the measurement period.

7.3 Data Characteristics

7.3.1 Parameter/Variable

The parameters contained in the data files on the CD-ROM are:

Column Name
SITE_NAME
SUB_SITE
DATE OBS
TIME OBS
PLOT_ID
WEIGHT
MEASUREMENT_TYPE
CRTFCN_CODE
REVISION DATE

7.3.2 Variable Description/Definition

The descriptions of the parameters contained in the data files on the CD-ROM are:

Column Name	Description
SITE_NAME	The identifier assigned to the site by BOREAS, in the format SSS-TTT-CCCCC, where SSS identifies the portion of the study area: NSA, SSA, REG, TRN, and TTT identifies the cover type for the site, 999 if unknown, and CCCCC is the identifier for site, exactly what it means will vary with site type.
SUB_SITE	The identifier assigned to the sub-site by BOREAS, in the format GGGGG-IIIII, where GGGGG is the group associated with the sub-site instrument, e.g. HYD06 or STAFF, and IIIII is the identifier for sub-site, often this will refer to an instrument.
DATE OBS	The date on which the data were collected.
TIME_OBS	The Greenwich Mean Time (GMT) when the data were collected.
PLOT_ID	The identifier for the plot from which the measurement came.

The following is needed to decode the PLOT_ID value. The characters indicate the layer(s) of moss that are in the turf and whether the turf had plastic applied to it or not.

L : The litter layer is included. F : The fibric layer is included. H : The humus layer is included.

: A number indicating the plot number of the turf.

p : plastic was applied, restricting the flow of water out of the turf.

WEIGHT
MEASUREMENT TYPE

The weight of the sample (including tray). Indicates whether the measurement was a "WET" measurement during the course of the field season or a "DRY" measurement after the field season

was over.

The information regarding MEASUREMENT_TYPE indicates whether the turf is WET or DRY. There is only one DRY measurement per turf, since the dry measurement was made when the turf was removed from the field, dried, and weighed in a lab.

CRTFCN_CODE

The BOREAS certification level of the data.

Examples are CPI (Checked by PI), CGR (Certified by Group), PRE (Preliminary), and CPI-??? (CPI but questionable).

REVISION_DATE

The most recent date when the information in the

referenced data base table record was revised.

7.3.3 Unit of Measurement

The measurement units for the parameters contained in the data files on the CD-ROM are:

Column Name	Units		
SITE NAME	[none]		
SUB SITE	[none]		
DATE_OBS	[DD-MON-YY]		
TIME_OBS	[HHMM GMT]		
PLOT_ID	[none]		
WEIGHT	[grams]		
MEASUREMENT_TYPE	[none]		
CRTFCN_CODE	[none]		
REVISION_DATE	[DD-MON-YY]		

7.3.4 Data Source

The sources of the parameter values contained in the data files on the CD-ROM are:

Column Name	Data Source
SITE_NAME	[Assigned by BORIS]
SUB_SITE	[Assigned by BORIS]
DATE OBS	[Supplied by Investigator]
TIME OBS	[Supplied by Investigator]
PLOT ID	[Supplied by Investigator]
WEIGHT	[Supplied by Investigator]
MEASUREMENT TYPE	[Supplied by Investigator]
CRTFCN CODE	[Assigned by BORIS]
REVISION_DATE	[Assigned by BORIS]

7.3.5 Data Range

The following table gives information about the parameter values found in the data files on the CD-ROM.

	Minimum	Maximum	Missng	Unrel	Below	Data
	Data	Data	Data	Data	Detect	Not
Column Name	Value	Value	Value	Value	Limit	Cllctd
SITE_NAME	NSA-9BS-HYD08	NSA-OBS-FLXTR	None	None	None	None
SUB_SITE	HYD08-MOS01	HYD08-MOS02	None	None	None	None
DATE_OBS	25-JUN-94	16-SEP-94	None	None	None	None
TIME_OBS	18	2309	None	None	None	None
PLOT_ID	L1	LFH4p	None	None	None	None
WEIGHT	18.07	1909.21	-99.9	None	None	None
MEASUREMENT_TYPE	DRY	WET	None	None	None	None
CRTFCN CODE	CPI	CPI	None	None	None	None
REVISION_DATE	07-JUL-97	10-JUL-97	None	None	None	None

Minimum Data Value -- The minimum value found in the column.

Maximum Data Value -- The maximum value found in the column.

Missng Data Value -- The value that indicates missing data. This is used to indicate that an attempt was made to determine the parameter value, but the attempt was unsuccessful.

Unrel Data Value -- The value that indicates unreliable data. This is used to indicate an attempt was made to determine the parameter value, but the value was deemed to be

unreliable by the analysis personnel.

Below Detect Limit -- The value that indicates parameter values below the instruments detection limits. This is used to

indicate that an attempt was made to determine the parameter value, but the analysis personnel determined that the parameter value was below the detection

limit of the instrumentation.

Data Not Clictd -- This value indicates that no attempt was made to

determine the parameter value. This usually indicates that BORIS combined several similar but not identical data sets into the same data base table

but this particular science team did not

measure that parameter.

Blank -- Indicates that blank spaces are used to denote that type of value.

N/A -- Indicates that the value is not applicable to the respective column.

-- Indicates that no values of that sort were found in the column.

7.4 Sample Data Record

The following are wrapped versions of data records from a sample data file on the CD-ROM.

SITE NAME, SUB SITE, DATE OBS, TIME OBS, PLOT ID, WEIGHT, MEASUREMENT TYPE, CRTFCN CODE, REVISION DATE

^{&#}x27;NSA-9BS-HYD08', 'HYD08-MOS01', 25-JUN-94, 1800, 'L1p', 141.5, 'WET', 'CPI', 10-JUL-97

^{&#}x27;NSA-OBS-FLXTR','HYD08-MOS02',15-AUG-94,2100,'L2',111.4,'WET','CPI',07-JUL-97

8. Data Organization

8.1 Data Granularity

The smallest amount of data that can be ordered is all of the data for a given site.

8.2 Data Format(s)

The Compact Disk-Read-Only Memory (CD-ROM) files contain American Standard Code for Information Interchange (ASCII) numerical and character fields of varying length separated by commas. The character fields are enclosed with single apostrophe marks. There are no spaces between the fields.

Each data file on the CD-ROM has four header lines of Hyper-Text Markup Language (HTML) code at the top. When viewed with a Web browser, this code displays header information (data set title, location, date, acknowledgments, etc.) and a series of HTML links to associated data files and related data sets. Line 5 of each data file is a list of the column names, and line 6 and following lines contain the actual data.

9. Data Manipulations

9.1 Formulae

None.

9.1.1 Derivation Techniques and Algorithms

The mass of water in a lysimeter can be estimated from this data set with the following equation:

```
mass_water(g) = wet_mass(g) - dry_mass(g) - tray_mass(g)
```

The computation of water equivalent depth for the lysimeters was performed using:

```
d (mm) = 1000 (mm/m) * mass_water(g) / (1000 kg/m<sup>3</sup> * area_gauge_bottom(m<sup>2</sup>))
```

9.2 Data Processing Sequence

9.2.1 Processing Steps

• Set up necessary equipment.

Performed daily weighings and empty weighed gauges.

Performed the necessary data manipulations and computed water equivalent depth.

Added the necessary column headings.

Transferred the information to the BOREAS Information System (BORIS).

Loaded the data into the relational data base (done by BORIS staff).

9.2.2 Processing Changes

None.

9.3 Calculations

9.3.1 Special Corrections/Adjustments

None given.

9.3.2 Calculated Variables

None given.

9.4 Graphs and Plots

None.

10. Errors

10.1 Sources of Error

Sampling error: errors might have occurred because of a limited number of samples.

10.2 Quality Assessment

10.2.1 Data Validation by Source

None given.

10.2.2 Confidence Level/Accuracy Judgment

None given.

10.2.3 Measurement Error for Parameters

None given.

10.2.4 Additional Quality Assessments

None.

10.2.5 Data Verification by Data Center

The data were received from the HYD-08 science team and loaded into the BORIS relational data base. After loading, the data were compared with the original data files to make sure that they were loaded properly.

11. Notes

11.1 Limitations of the Data

None given.

11.2 Known Problems with the Data

None given.

11.3 Usage Guidance

The weights that are reported in these data can be used to compute equivalent water depth. Equivalent water depth can be computed by using dry turf + lysimeter tray weights to compute water weight, assuming a constant 1,000 kg/m³ density of H_2O to compute water equivalent depth for a 20-cm x 20-cm turf sample.

11.4 Other Relevant Information

The tray weights are included in the weights reported in this data set. When comparing wet and dry turf weights, the weight of the tray cancels out to yield the weight of the water.

12. Application of the Data Set

The data collected may be useful in characterizing canopy interception, drip, throughfall, moss interception, drainage, evaporation, and capacity during the growing season at daily temporal resolution.

13. Future Modifications and Plans

None given.

14. Software

14.1 Software Description

None.

14.2 Software Access

Not applicable.

15. Data Access

The HYD-08 1994 gravimetric moss moisture data are available from the Earth Observing System Data and Information System (EOSDIS) Oak Ridge National Laboratory (ORNL) Distributed Active Archive Center (DAAC).

15.1 Contact Information

For BOREAS data and documentation please contact:

ORNL DAAC User Services Oak Ridge National Laboratory P.O. Box 2008 MS-6407 Oak Ridge, TN 37831-6407 Phone: (423) 241-3952

Fax: (423) 574-4665

E-mail: ornldaac@ornl.gov or ornl@eos.nasa.gov

15.2 Data Center Identification

Earth Observing System Data and Information System (EOSDIS) Oak Ridge National Laboratory (ORNL) Distributed Active Archive Center (DAAC) for Biogeochemical Dynamics http://www-eosdis.ornl.gov/.

15.3 Procedures for Obtaining Data

Users may obtain data directly through the ORNL DAAC online search and order system [http://www-eosdis.ornl.gov/] and the anonymous FTP site [ftp://www-eosdis.ornl.gov/data/] or by contacting User Services by electronic mail, telephone, fax, letter, or personal visit using the contact information in Section 15.1.

15.4 Data Center Status/Plans

The ORNL DAAC is the primary source for BOREAS field measurement, image, GIS, and hardcopy data products. The BOREAS CD-ROM and data referenced or listed in inventories on the CD-ROM are available from the ORNL DAAC.

16. Output Products and Availability

16.1 Tape Products

None.

16.2 Film Products

None.

16.3 Other Products

These data are available on the BOREAS CD-ROM series.

17. References

17.1 Platform/Sensor/Instrument/Data Processing Documentation None.

17.2 Journal Articles and Study Reports

Haddeland, I. and D.P. Lettenmaier. 1995. Hydrologic Modeling of Boreal Forest Ecosystems. Water Resources Series Technical Report No. 143. University of Washington, 123 pp.

Newcomer, J., D. Landis, S. Conrad, S. Curd, K. Huemmrich, D. Knapp, A. Morrell, J. Nickeson, A. Papagno, D. Rinker, R. Strub, T. Twine, F. Hall, and P. Sellers, eds. 2000. Collected Data of The Boreal Ecosystem-Atmosphere Study. NASA. CD-ROM.

Price, A.G., K. Dunham, T. Carleton, and L.E. Band. 1997. Variability of water fluxes through the Black Spruce (Picea Mariana) canopy and Feather Moss (Pleurozium Schreberi) carpet in the Boreal Forest of Northern Manitoba. Journal of Hydrology, 196, 310-323.

Sellers, P. and F. Hall. 1994. Boreal Ecosystem-Atmosphere Study: Experiment Plan. Version 1994-3.0, NASA BOREAS Report (EXPLAN 94).

Sellers, P. and F. Hall. 1996. Boreal Ecosystem-Atmosphere Study: Experiment Plan. Version 1996-2.0, NASA BOREAS Report (EXPLAN 96).

Sellers, P., F. Hall, and K.F. Huemmrich. 1996. Boreal Ecosystem-Atmosphere Study: 1994 Operations. NASA BOREAS Report (OPS DOC 94).

Sellers, P., F. Hall, and K.F. Huemmrich. 1997. Boreal Ecosystem-Atmosphere Study: 1996 Operations. NASA BOREAS Report (OPS DOC 96).

Sellers, P., F. Hall, H. Margolis, B. Kelly, D. Baldocchi, G. den Hartog, J. Cihlar, M.G. Ryan, B. Goodison, P. Crill, K.J. Ranson, D. Lettenmaier, and D.E. Wickland. 1995. The boreal ecosystem-atmosphere study (BOREAS): an overview and early results from the 1994 field year. Bulletin of the American Meteorological Society. 76(9):1549-1577.

Sellers, P.J., F.G. Hall, R.D. Kelly, A. Black, D. Baldocchi, J. Berry, M. Ryan, K.J. Ranson, P.M. Crill, D.P. Lettenmaier, H. Margolis, J. Cihlar, J. Newcomer, D. Fitzjarrald, P.G. Jarvis, S.T. Gower, D. Halliwell, D. Williams, B. Goodison, D.E. Wickland, and F.E. Guertin. 1997. BOREAS in 1997: Experiment Overview, Scientific Results and Future Directions. Journal of Geophysical Research 102(D24): 28,731-28,770.

17.3 Archive/DBMS Usage Documentation None.

18. Glossary of Terms

None.

ASCII

19. List of Acronyms

BOREAS - BOReal Ecosystem-Atmosphere Study
BORIS - BOREAS Information System
CD-ROM - Compact Disk-Read-Only Memory
DAAC - Distributed Active Archive Center
EOS - Earth Observing System
EOSDIS - EOS Data and Information System

- American Standard Code for Information Interchange

FFC-T - Focused Field Campaign - Thaw
GIS - Geographic Information System

GMT - Greenwich Mean Time

GSFC - Goddard Space Flight Center HTML - Hyper-Text Markup Language

HYD - Hydrology

IFC - Intensive Field Campaign

MSA - Modeling Sub-Area

NAD83 - North American Datum of 1983

NASA - National Aeronautics and Space Administration

NSA - Northern Study Area

OBS - Old Black Spruce Tower Site
ORNL - Oak Ridge National Laboratory
PANP - Prince Albert National Park

SSA - Southern Study Area
URL - Uniform Resource Locator

WWW - World Wide Web

20. Document Information

20.1 Document Revision Date

Written: 15-Nov-1995 Last Revised: 16-Jul-1999

20.2 Document Review Date(s)

BORIS Review: 23-Jul-1998

Science Review:

20.3 Document ID

20.4 Citation

When using these data, please contact the principal investigator, Dr. Lawrence Band (see Section 2.1), before publishing results that are based on these data as well as citing relevant papers in Section 17.2.

If using data from the BOREAS CD-ROM series, also reference the data as:

Band, L.E., "Simulation of Boreal Ecosystem Carbon and Water Budgets: Scaling from Local to Regional Extents." In Collected Data of The Boreal Ecosystem-Atmosphere Study. Eds. J. Newcomer, D. Landis, S. Conrad, S. Curd, K. Huemmrich, D. Knapp, A. Morrell, J. Nickeson, A. Papagno, D. Rinker, R. Strub, T. Twine, F. Hall, and P. Sellers. CD-ROM. NASA, 2000.

Also, cite the BOREAS CD-ROM set as:

Newcomer, J., D. Landis, S. Conrad, S. Curd, K. Huemmrich, D. Knapp, A. Morrell, J. Nickeson, A. Papagno, D. Rinker, R. Strub, T. Twine, F. Hall, and P. Sellers, eds. Collected Data of The Boreal Ecosystem-Atmosphere Study. NASA. CD-ROM. NASA, 2000.

20.5 Document Curator

20.6 Document URL

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington to Washington to Reports, 1215 Jefferson Davis Highery, Suite 1204 Adjusting VA 2202.4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)	GENCY USE ONLY (Leave blank) 2. REPORT DATE July 2000 3. REPORT TYPE AT Technical M		D DATES COVERED emorandum	
4. TITLE AND SUBTITLE	July 2000	Teenmeat W	5. FUNDING NUMBERS	
Technical Report Series on the Bo	oreal Ecosystem-Atmosphe	ere Study (BOREAS)		
BOREAS HYD-8 1994 Grav			923	
			RTOP: 923-462-33-01	
6. AUTHOR(S) Xuewen Wang				
Forrest G. Hall and David E.	Knapp Editors			
Tonest G. Han and David E.	Khapp, Editors			
7. PERFORMING ORGANIZATION NAM	E(S) AND ADDRESS (ES)		8. PEFORMING ORGANIZATION REPORT NUMBER	
Goddard Space Flight Center			The one nomber	
Greenbelt, Maryland 20771			2000-03136-0	
9. SPONSORING / MONITORING AGE	ENCY NAME(S) AND ADDRES	SS (ES)	10. SPONSORING / MONITORING AGENCY REPORT NUMBER	
National Aeronautics and Space	• Administration		TM-2000-209891	
Washington, DC 20546-0001	, rigininistration		Vol. 34	
,, asimigron, 2 0 200 to 0001				
			1	
11. SUPPLEMENTARY NOTES				
X. Wang: University of Toron	nto; D.E. Knapp: Raythe	eon ITSS		
12a. DISTRIBUTION / AVAILABILITY ST	ATEMENT	<u> </u>	12b. DISTRIBUTION CODE	
Unclassified-Unlimited				
Subject Category: 43				
Report available from the NAS.				
7121 Standard Drive, Hanover,	MD 21076-1320. (301) 6	521-0390.		
13. ABSTRACT (Maximum 200 words)				
	1	Co aa ladual	al processes that were collected	
The RODEAS HVI)-8 team:	made measurements of a	surtace nyaratogica	n processes inal were collecte	

The BOREAS HYD-8 team made measurements of surface hydrological processes that were collected at the NSA-OBS Tower Flux site in 1994 and at Joey Lake, Manitoba, to support its research into point hydrological processes and the spatial variation of these processes. The data collected may be useful in characterizing canopy interception, drip, throughfall, moss interception, drainage, evaporation, and capacity during the growing season at daily temporal resolution. This particular data set contains the gravimetric moss moisture measurements from June to September 1994. A nested spatial sampling plan was implemented to support research into spatial variations of the measured hydrological processes and ultimately the impact of these variations on modeled carbon and water budgets. These data are stored in tabular ASCII files.

14. SUBJECT TERMS BOREAS, hydrology, gra	15. NUMBER OF PAGES 14 16. PRICE CODE		
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT UL